

THE MAPLE[®] SYMBOLIC MATHEMATICS SYSTEM IN THE METHOD OF PROJECTIONS FOR DISCRETE OPTIMIZATION PROBLEMS**Chernova L.S.**

*Admiral Makarov National University of Shipbuilding
Ukraine, 54025, Mykolayiv, Prospekt Heroiv Ukrainy, 9
19chsk56@gmail.com, ss1-ss10@ukr.net*

ORCID ID 0000-0003-3954-7436

Key words:

IT project management, education programs aimed at Master's degree, linear optimization, discrete optimization, system of constraints, combinatory method, the Gauss-Jordan method, decomposition, reduction, graphical solution, Maple[®].

Successful project activities in the IT industry are determined by the extent of difficulty in the team formation and implementation of projects themselves. IT projects provide for the fulfilment of a number of tasks that are interrelated. In the formation of such a project, account should be taken of certain factors necessary for its successful implementation, determination of the technology for fulfilling intermediate tasks: consecutive or parallel, and for setting the priorities. This approach requires detailed calculation and scientifically grounded decisions. The authors have proposed an original approach to solving discrete optimization problems related to fundamental calculation difficulties in the process of an IT project formation. The known methods of exact or approximate solution of such problems are studied with account taken of their belonging to so-called P- and NP-class problems (the polynomial and the exponential solution algorithms). The modern combinatory and heuristic methods for solving practical discrete optimization problems require development of algorithms that allow obtaining approximate solution with guaranteed estimate of deviation from the optimum. Simplification algorithms provide an efficient method of searching for an optimization problem solution. Should a multidimensional process be projected onto the two-dimensional surface, this will enable graphical visualization of sets of the problem solutions. This research provides a way for simplifying the combinatory solution of a discrete optimization problem. It is based on decomposition of the system that represents the system constraining a multidimensional output problem to the two-dimensional coordinate plane. Such method allows obtaining a simple system of graphical solutions of a complicated linear discrete optimization problem. From the practical point of view, the proposed method allows reducing the calculation compliance of optimization problems belonging to this class when the IT project solutions are complicated. The approach proposed can be applied in using the obtained research result for assuring the possibility to improve the class of problems presented by linear equation systems. The automation of calculations in the Maple[®] environment provides the basis for further development and improvement of such algorithms, and for using in teaching a number of disciplines in education programs on IT project management aimed at Master's degree.

СИСТЕМА СИМВОЛЬНОЇ МАТЕМАТИКИ MAPLE[®] У МЕТОДІ ПРОЄКЦІЙ ДЛЯ ЗАДАЧ ДИСКРЕТНОЇ ОПТИМІЗАЦІЇ**Чернова Л.С.**

*Національний університет кораблебудування ім. Адмірала Макарова
Україна, 54025, м. Миколаїв, пр. Героїв України, 9*

Ключові слова:

управління ІТ проєктами, освітні програми магістерського освітньо-кваліфікаційного рівня, лінійна оптимізація, дискретна оптимізація, система обмежень, комбінаторний метод, метод Жордана – Гаусса, декомпозиція, редукція, графічний розв'язок, Maple[®].

Успішна проєктна діяльність у ІТ галузі визначається складністю формування команди та реалізації самих проєктів. ІТ проєкти передбачають виконання ряду взаємопов'язаних завдань. При формуванні такого проєкту слід враховувати низку чинників, необхідних для його успішної реалізації, визначення технології реалізації проміжкових завдань: послідовного паралельного і завдання пріоритетності. Такий підхід потребує детальних розрахунків та науковообґрунтованих рішень. Автори пропонують оригінальний підхід до розв'язку задач дискретної оптимізації, пов'язаних із принциповими обчислювальними труднощами в процесі формування ІТ проєкту. Відомі методи точного або наближеного розв'язку таких задач вивчаються з урахуванням належності їх до, так званих, задач з класу P та NP (алгоритми поліноміальної та експоненціальної реалізації розв'язку). Сучасні комбінаторні та евристичні методи розв'язку практичних задач дискретної оптимізації потребують розробки алгоритмів, які дозволяють отримувати наближений розв'язок із гарантованою оцінкою відхилення від оптимуму. Алгоритми спрощення є ефективним прийомом пошуку розв'язку оптимізаційної задачі. Якщо виконати проєктування багатовимірною процесу на двовимірну площину, то такий прийом

дозволить наочно відобразити в графічній формі множини розв'язків задачі. У межах дослідження запропоновано спосіб спрощення комбінаторного розв'язку задач дискретної оптимізації. Він заснований на тому, що виконується декомпозиція системи, яка відображає систему обмежень багатовимірної вихідної задачі на двовимірну координатну площину. Такий спосіб дозволяє отримати просту систему графічних розв'язувань складної задачі лінійної дискретної оптимізації. Із практичної точки зору запропонований метод дозволяє спростити обчислювальну складність оптимізаційних задач такого класу при складних ІТ проектних рішеннях. Прикладним аспектом запропонованого підходу є використання отриманого наукового результату для забезпечення можливості вдосконалення такого класу задач, що описуються системами лінійних рівнянь. Автоматизація розрахунків у середовищі Maple® створює передумови для подальшого розвитку та удосконалення подібних алгоритмів та використання для викладання ряду предметів в освітніх програмах з управління ІТ проектами магістерського освітньо-кваліфікаційного рівня.

Statement of the problem

Discrete optimization problems appear in many areas where models of current processes were formed and use mathematical methods of their solution under the following additional conditions: the unknowns have to be integer in full or in part, or they have to be binary (0 or 1). The travelling salesman problem, the knapsack problem and the assignment problem are the most known problems of linear optimization. Nowadays, discrete optimization has been formed as an independent part of the theory of optimization. It uses modern combinatorial methods and algorithms for solving practical problems. Their application results in primary bases of a problem, further assessment of their optimality, improvement of bases in case of their nonoptimality and bounds of the target function [1,2,3,4].

As the most discrete optimization problems belong to the *NP* class, it is reasonable to use problem simplification algorithms without losing the controlled accuracy of solution [5,6,7]. The procedure of simplification uses the known interrelation of linear algebraic equation systems with the system of linear algebraic inequalities and the classic linear algebra apparatus [8,9,10].

The principle of the method proposed consists in using the feature of convex polyhedral area Ω_I provided by a system of linear algebraic inequalities or equations in the form of a direct sum of subspace and kernel [11]. Provided that the polyhedral kernel is two-dimensional, an optimization problem can be reduced to a two-dimensional one. Projections obtained enable easy finding of the optimum solution and evaluation of availability of an integer solution, and then of a binary solution as well. A direct calculation means of simplifying such class of discrete optimization problems that are implemented by the Gauss-Jordan method in the Maple® computer mathematics environment [7,10,12,13].

Analysis of recent studies and publications

Mathematical models of active systems are interpreted in many cases as discrete optimization problems [1,2,14,15]. Development of discrete optimization problems is associated with fundamental difficulties [2]. The known modern methods and algorithms of exact and approximate solution of such problems are studied

with account taken of their belonging to so-called *P*- and *NP*-class problems (the polynomial and the exponential solution algorithms) [5].

Combinatorial and heuristic methods for exact and approximate solving practical discrete optimization problems take an essential place in obtaining optimum values of such problems [1]. Realization of such algorithms requires availability of the acceptable primary basis of a problem, an optimality assessment procedure and the basis improvement if nonoptimality is the case [5,6].

The methods of discrete optimization problems solution that have been developed by now require development of algorithms which allow obtaining an approximate solution with guaranteed estimate of deviation from the optimum.

Simplification algorithms in discrete optimization problems provide an efficient method of searching for an optimization problem solution [16,17,18]. Should a multidimensional process be projected onto the two-dimensional surface, this will enable visualization of the acceptable set (array) of the problem parameters. We can make a lower- and an upper-bound estimate of the problem target function values and dynamically evaluate the possibility to diversify basis optimum variables with guaranteed accuracy.

Solving contradictions between requirements to completeness of modelling views in active systems and methods of obtaining solutions of their mathematical models is possible due to reasonable reduction of the algorithms of complicated equation systems solution [17]. Lack of problem solution as regards searching for solutions in discrete optimization problems consists in the need for developing and implementing the procedure of simplifying the combinatorial solution of a discrete optimization problem.

Objectives of the article

The following objectives are determined for the research: involving and using linear algebra standard calculation procedures and certain linear optimization methods to simplify the solution of multidimensional discrete optimization problems with further visualizing of the geometric interpretation of a linear discrete optimization problem solution.

The following tasks were set to achieve the objectives determined:

- to remove the class of problems that have to be

simplified;

- to provide calculations of a modelling example.

The main material of the research

Development of simplification of the solution in discrete optimization problems

Successful project activities in the IT industry are determined by the extent of difficulty in the team formation and implementation of projects themselves. IT projects provide for the fulfilment of a number of tasks that are interrelated. In the formation of such a project, account should be taken of certain factors necessary for its successful implementation, determination of the technology for fulfilling intermediate tasks: consecutive or parallel, and for setting the priorities. To fulfil the task of a project content optimization, works are analysed that have to be performed for creating a software. In this process, the project is analysed for its content, period for each stage implementation, costs, risks and value. The value determining approach is based on comprehensive characteristic of the project results. This characteristic, in its turn, can be defined by

quality of the software created as a result of the project implementation, as well as by economic, social and politic, environmental, technical and other effects. For efficient calculations and avoiding fundamental calculation difficulties in the process of an IT project formation, evaluation of project actions in all phases of the project lifecycle – from issuance of requirements specifications to software installation to the customer, the authors have developed a method reducing the calculation compliance of optimization problems.

This research proposed the system decomposition through projection of a multidimensional output problem onto two-dimensional coordinate planes. This method transforms the output problem into a group of subsystems, which enables obtaining the system of graphical solutions of a complicated linear discrete optimization problem. The Maple® software environment has been involved from the methodological and scientific points of view. Each of the problem stages has been associated with a subprogram for automation of calculations and visualization of solution results (Fig. 1).




Fig. 1. Enlarged UML diagram of information system

For the method use visualization, we formulate a problem on the optimum placement of n - sets A_j ,

$j = 1, \dots, n$ on the universal set U . Let each set A_j , $j = 1, \dots, n$ be characterized by two scalar values: c_j

- value and a_j - power or weight $|A_j|$. At the same time, the condition is fulfilled that the power of universal

set $|U| = B$ is smaller than the total power of all A_j .

Such an optimization problem consists in the need to select a certain number of A_j from the total aggregate of A_j , for immersion into U , the total value of which is maximum.

The total power $\sum_{j=1}^n a_j > B$ is bigger than the power of universal set, i.e. it is impossible to place the complete number of sets. U can only accommodate a part (a number) of sets A_j . Let us enter n Boolean (dichotomic or binary) variables:

$$x_j = \begin{cases} 0, & A_j \text{ is not placed in } U, \\ 1, & A_j \text{ is placed in } U, \end{cases} \quad (1)$$

where $j = 1, \dots, n$.

Entering binary unknowns (1) - $x_j = 0 \vee 1$,

$j = 1, \dots, n$ allows formulating the target function

W_I and constraint Ω_I of the following optimization problem:

$$W_I = c_1x_1 + c_2x_2 + \dots + c_nx_n \rightarrow \max,$$

$$\Omega_I : a_1x_1 + a_2x_2 + \dots + a_nx_n \leq B, \quad (2)$$

$$x_j = 0 \vee 1, \quad j = 1, 2, \dots, n.$$

According to the problem statement, $c_j > 0$,

$$0 < a_j \leq B, \quad j = 1, 2, \dots, n.$$

Such problem (2) is called in the discrete optimization theory a one-dimensional knapsack problem. The solution of this problem means finding among 2^n n -dimensional vectors such vector

$$X = [x_1, x_2, \dots, x_n],$$

which meets the constraints Ω_I and provides the maximum value to the target

function W_I (Fig. 2).




Fig. 2. The problem on A_j immersion into U . (one-dimensional knapsack)

Let us consider the generalized one-dimensional knapsack problem statement. We divide the universal set into its own subsets $U = U_1 \cup U_2 \cup \dots \cup U_m$ with the condition $|U_i| = b_i, \quad i = 1, 2, \dots, m$, and $B = \sum_{i=1}^m b_i$. The problem of immersion of sets $A_j, \quad j = 1, \dots, n$ into $U = U_1 \cup U_2 \cup \dots \cup U_m$ is interpreting A_j as a set not just with one feature a_j , but with a whole range of $a_{ij}, \quad i = 1, 2, \dots, m$. The features are provided by set $A = [a_{ij}]_{m \times n}$. The mathematic form of such optimization problem on placement of A_j into $U = U_1 \cup U_2 \cup \dots \cup U_m$ looks as follows:

$$W_I = c_1x_1 + c_2x_2 + \dots + c_nx_n \rightarrow \max,$$

$$\Omega_I : \begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n \leq b_1, \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n \leq b_2, \\ \vdots \quad \dots \quad \dots \quad \vdots \\ a_{m-1,1}x_1 + a_{m-1,2}x_2 + \dots + a_{m-1,n}x_n \leq b_{m-1}, \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n \leq b_m, \end{cases} \quad (3)$$

$$x_j = 0 \vee 1, \quad j = 1, 2, \dots, n.$$

Account is to be taken of the fact that $c_j > 0$,

$$0 < a_{ij} \leq B, \quad i = 1, 2, \dots, m, \quad j = 1, 2, \dots, n.$$

For $\forall i \in [1, m]$ the following is to fulfil:

$$a_{i1} + a_{i2} + \dots + a_{in} > b_i.$$

It means that it is not possible to place all sets $A_j, \quad j = 1, \dots, n$ in any of the

subsets $U = U_1 \cup U_2 \cup \dots \cup U_m$. The number of n – measurable vectors $X = [x_1, x_2, \dots, x_n]$ is the solution of the problem.

Such type of problem is used to be called a one-dimensional knapsack problem. The formulated problem is interpreted as a problem on the optimum selection in project management.

For implementation of n projects ($A_j, j = 1, \dots, n$), certain resources are provided that are represented in the form of a vector of resources $b = [b_1, b_2, \dots, b_m]^T$

The set $A = [a_{ij}]_{m \times n}$ determines the rates of consuming resource b_j for implementation of project A_j . The profit from implementation of project A_j is $c_j > 0$. We need to choose the number of projects A_j that allows gaining the maximum profit. Let us enter Boolean vector $X = [x_1, x_2, \dots, x_n]$, where

$$x_j = \begin{cases} 0, & \text{project } A_j \text{ is not implemented,} \\ 1, & \text{project } A_j \text{ implemented,} \end{cases} \\ j = 1, \dots, n.$$

We obtain the problem:

$$W_I = CX^T \rightarrow \max, \\ \Omega_I : AX \leq b, \tag{4}$$

$$x_j = 0 \vee 1, \quad j = 1, 2, \dots, n.$$

The project management model (4) completely agrees with the multidimensional knapsack problem (3).

At this time, we know that the given problems (3), (4) cannot be exactly resolved. We have performed exhaustive research of features of acceptable and optimum solutions of knapsack problems and proposed several algorithms [2,5] of gradual approximating to the optimum solution. Thus, the Danzig algorithm and so-called “greedy” procedures form the basis of heuristic algorithms [18].

The research has proposed and exactly grounded an approach to finding the optimum solution of a broad class of multidimensional knapsack problems. The principle of the method proposed consists in using the feature of convex polyhedral area Ω_I provided by a system of linear algebraic inequalities or equations in the form of a direct sum of subspace and kernel [10, 17]. Provided that the polyhedral kernel is two-dimensional, an optimization problem can be reduced to a two-dimensional one. Projections obtained enable easy finding of the optimum solution and evaluation of availability of an integer solution, and then of a binary solution as well.

In other words, the research proposed projecting polyhedron Ω_I onto subsets of the set of basis vectors of a linear optimization problem system of constraints.

For special case $m - n \leq 2$, where m - number of constraints Ω_I , n - rank $A = [a_{ij}]_{m \times n}$ projections are elementary because there are two-dimensional. It is not difficult to analyse the projection integer values array and to solve the problem.

The first step in this algorithm is to prepare the system of constraints for reduction. Thus, let us have a general optimization problem in the following form:

$$W_I = \sum_{j=1}^n c_j x_j \rightarrow \max, \\ \Omega_I : \begin{cases} \sum_{j=1}^n a_{ij} x_j \leq b_i, & i = 1, \dots, k, \\ \sum_{j=1}^n a_{ij} x_j = b_i, & i = k + 1, \dots, m, \end{cases} \tag{3} \\ x_j \geq 0, \quad j = 1, \dots, l.$$

We know that such a problem can be reduced to the canonical form:

$$W_I = \sum_{j=1}^n c_j x_j \rightarrow \max, \\ \Omega_I : \sum_{j=1}^n a_{ij} x_j = b_i, \quad i = 1, \dots, m, \\ x_j \geq 0, \quad j = 1, \dots, n.$$

The reduction is possible due to standard methods of transformation. Thus, the equation of system of constraints is equivalent to the system of two inequalities:

$$\sum_{j=1}^n a_{ij} x_j = b_i \Leftrightarrow \begin{cases} \sum_{j=1}^n a_{ij} x_j \leq b_i, \\ -\sum_{j=1}^n a_{ij} x_j \leq -b_i. \end{cases}$$

Values with arbitrary sign can be represented in the form of a difference of 2 nonnegative variables:

$$x_j = u_j - v_j, \quad u_j \geq 0, \quad v_j \geq 0.$$

Transition from inequality constraints to equation constraints is made by adding the nonnegative (balancing) variable:

$$\sum_{j=1}^n a_{ij} x_j \leq b_j \Rightarrow \sum_{j=1}^n a_{ij} x_j + x_{n+i} = b_i, \\ x_{n+i} \geq 0, \quad i = 1, \dots, k.$$

Transition from maximization to minimization of the target function and transition the other way round is used for simpler transformation:

$$W_I = \sum_{j=1}^n c_j x_j \rightarrow \max \Leftrightarrow W_I = -\sum_{j=1}^n c_j x_j \\ \rightarrow \min$$

In view of this, without loss of considerations generality, let us have a linear optimization problem provided in canonical form:

$$W_I = CX \rightarrow \max$$

$$\Omega_I : AX = B,$$

$$X \geq 0,$$

where the matrix rank of factors of the system of constraints is equal to rank $A = m$.

Solving the system with the Gauss-Jordan method under an arbitrary basis combination of variables, we obtain projection n - of measurable output problem onto $(m - n)$ - measurable space. As we take into consideration a class of problems with condition $m - n = 2$, we have projecting of R^n onto two-dimensional plane R^2 .

Let us consider a modelling example of solution based on projecting a multidimensional process in R^6 onto two-dimensional space R^2 .

Modelling example

Tasks: solving the optimization problem with condition $x_j \geq 0, j = 1, \dots, n$. Also obtaining a completely integer solution and the solution under condition $x_j = 0 \vee 1, j = 1, 2, \dots, n$.

$$W_I = x_1 + 8x_2 + 4x_3 + x_4 \rightarrow \max,$$

$$\Omega_I : \begin{cases} 2x_1 + 3x_2 + 4x_3 + 3x_4 \leq 7, \\ 3x_1 + 5x_2 + 2x_3 + 4x_4 = 8, \\ 3x_1 + 3x_2 + 5x_3 + 4x_4 = 6, \\ 3x_1 + 5x_2 + 3x_3 + 2x_4 \leq 8, \end{cases}$$

$$x_1 \geq 0, x_2 \geq 0, x_3 \geq 0, x_4 \geq 0.$$

Table 1

	x_1	x_2	x_3	x_4	x_5	x_6	b	Σ
	2	3	4	3	1	0	7	20
	3	5	2	4	0	0	8	22
	3	3	5	4	0	0	6	21
	3	5	3	2	0	1	8	22
W_I	1	8	4	1	0	0	0	
	0	- 1/3	8/3	1/3	1	0	5/3	16/3
	1	5/3	2/3	4/3	0	0	8/3	22/3
	0	- 2	3	0	0	0	- 2	- 1
	0	0	1	- 2	0	1	0	0
W_I	0	19/3	10/3	- 1/3	0	0	- 8/3	
	0	0	13/6	1/3	1	0	2	11/2
	1	0	19/6	4/3	0	0	1	13/2
	0	1	- 3/2	0	0	0	1	1/2
	0	0	1	- 2	0	1	0	0
W_I	0	0	77/6	- 1/3	0	0	- 9	

From the last transformation of Table 1, we have the solved system

We make a transition to canonical form of a linear optimization problem

$$W_I = x_1 + 8x_2 + 4x_3 + x_4 \rightarrow \max,$$

$$\Omega_I : \begin{cases} 2x_1 + 3x_2 + 4x_3 + 3x_4 + x_5 = 7, \\ 3x_1 + 5x_2 + 2x_3 + 4x_4 = 8, \\ 3x_1 + 3x_2 + 5x_3 + 4x_4 = 6, \\ 3x_1 + 5x_2 + 3x_3 + 2x_4 + x_6 = 8, \end{cases}$$

$$x_j \geq 0, j = 1, \dots, 6.$$

The system of constraints consists of four independent equations $rank(A) = 4$. We move from the canonical form of problem representation to the standard form. Such move (projecting) is made with solving the system by the Gauss-Jordan method. (Table 1) For the given problem of projection $R^6 \Rightarrow R^2$ we can perform $C_6^4 = 15$ ways. We choose randomized basis combination Ox_3x_4 .

Projection onto Ox_3x_4

As basis variables, we choose the following quadruple: x_1, x_2, x_5, x_6 .

$$\begin{aligned}
 & W_l = x_1 + 8x_2 + 4x_3 + x_4 \rightarrow \max, \\
 \Omega_l : & \begin{cases} \frac{13}{6}x_3 + \frac{1}{3}x_4 + x_5 = 2, \\ \frac{19}{6}x_3 + \frac{4}{3}x_4 + x_1 = 1, \\ -\frac{3}{2}x_3 + x_2 = 1, \\ x_3 - 2x_4 + x_6 = 0, \\ x_j \geq 0, \quad j = 1, \dots, 6. \end{cases} \tag{5}
 \end{aligned}$$

Truncating the basis variables, we assure transition $R^6 \Rightarrow R^2$ to two-dimensional inequalities. The projection of six-dimensional output problem onto

coordinate plane Ox_3x_4 has the following analytical form:

$$\begin{aligned}
 & W_l = \frac{77}{6}x_3 - \frac{1}{3}x_4 + 9 \rightarrow \max, \\
 \Omega_l^{Ox_3x_4} : & \begin{cases} \frac{13}{6}x_3 + \frac{1}{3}x_4 \leq 2, \\ \frac{19}{6}x_3 + \frac{4}{3}x_4 \leq 1, \\ -\frac{3}{2}x_3 \leq 1, \\ x_3 - 2x_4 \leq 0, \\ x_3 \geq 0, \quad x_4 \geq 0, \end{cases} \Leftrightarrow \Omega_l^{Ox_3x_4} : \begin{cases} 13x_3 + 2x_4 \leq 12, \\ 19x_3 + 8x_4 \leq 6, \\ -3x_3 \leq 2, \\ x_3 - 2x_4 \leq 0, \\ x_3 \geq 0, \quad x_4 \geq 0. \end{cases}
 \end{aligned}$$

The graphical solution is given on Fig. 3.




Fig. 3. Projection onto Ox_3x_4

The solution of the system is provided in the extremal vertex coordinates

$$X_{\max}^{opt} : \omega_2 \times \omega_4 \Leftrightarrow \begin{cases} 19x_3 + 8x_4 = 6, \\ x_3 - 2x_4 = 0, \end{cases} \Leftrightarrow \begin{cases} x_3 = \frac{6}{23}, \\ x_4 = \frac{3}{23}. \end{cases}$$

Other coordinates are to obtain from the solved system (5) Therefore, the optimum solution of the output problem is equal to:

$$X_{\max}^{opt} = \left[0, \frac{32}{23}, \frac{6}{23}, \frac{3}{23}, \frac{32}{23}, 0 \right].$$

The biggest value of the target function is $W_I^{\max} = \frac{289}{23}$.

The represented $\Omega_I^{Ox_3x_4}$ projection Ω_I onto Ox_3x_4 (Fig. 2) lets us state that point (0,0) is the only integer solution point of the problem. In view of this, we have the first integer optimum solution estimate $X_{\max}^Z = [x_1, x_2, 0, 0,]$.

Projection onto Ox_1x_2

For calculation of x_1, x_2 values, we project Ω_I onto Ox_1x_2 .

As basis variables, we choose the following quadruple: x_3, x_4, x_5, x_6 . We make calculations with the Gauss-Jordan method. (Table 2).

Table 2

	x_1	x_2	x_3	x_4	x_5	x_6	b	Σ
	2	3	4	3	1	0	7	20
	3	5	2	4	0	0	8	22
	3	3	5	4	0	0	6	21
	3	5	3	2	0	1	8	22
W_I	1	8	4	1	0	0	0	
	-4	-7	0	-5	1	0	-9	-24
	1,5	2,5	1	2	0	0	4	11
	-4,5	-9,5	0	-6	0	0	-14	-34
	-1,5	-2,5	0	-4	0	1	-4	-11
W_I	-5	-2	0	-7	0	0	-16	
	- 1/4	11/12	0	0	1	0	8/3	13/3
	0	- 2/3	1	0	0	0	- 2/3	- 1/3
	3/4	19/12	0	1	0	0	7/3	17/3
	3/2	23/6	0	0	0	1	16/3	35/3
W_I	1/4	109/12	0	0	0	0	1/3	

From the last transformation of Table 2, we have the solved system

$$\begin{aligned}
 &W_I = x_1 + 8x_2 + 4x_3 + x_4 \rightarrow \max, \\
 &\Omega_I : \begin{cases} -\frac{1}{4}x_1 + \frac{11}{12}x_2 + x_5 = \frac{8}{3}, \\ -\frac{2}{3}x_2 + x_3 = -\frac{2}{3}, \\ \frac{3}{4}x_1 + \frac{19}{12}x_2 + x_4 = \frac{7}{3}, \\ \frac{3}{2}x_1 + \frac{23}{6}x_2 + x_6 = \frac{16}{3}, \\ x_j \geq 0, \quad j = 1, \dots, 6. \end{cases} \tag{6}
 \end{aligned}$$

It should be mentioned that it is already enough just to have this only system for finding an integer solution. Indeed, we ascertained from the previous projecting that $x_3 = 0$ and $x_4 = 0$. In view of this, the second system equation gives $x_2 = 1$, and the third equation gives $x_1 = 1$. Therefore, $X_{\max}^Z = [1, 1, 0, 0,]$ is the

$$W_f = \frac{1}{4}x_1 + \frac{109}{12}x_2 - \frac{1}{3} \rightarrow \max,$$

$$\Omega_f^{Ox_1x_2} : \begin{cases} -\frac{1}{4}x_1 + \frac{11}{12}x_2 \leq \frac{8}{3}, \\ -\frac{2}{3}x_2 \leq -\frac{2}{3}, \\ \frac{3}{4}x_1 + \frac{19}{12}x_2 \leq \frac{7}{3}, \\ \frac{3}{2}x_1 + \frac{23}{6}x_2 \leq \frac{16}{3}, \\ x_1 \geq 0, x_2 \geq 0. \end{cases}$$

integer solution of the problem. We confirm these values through the graphical solution of the problem. Truncating the basis variables, we assure transition $R^6 \Rightarrow R^2$ to two-dimensional inequalities. The projection of six-dimensional output problem onto coordinate plane Ox_1x_2 has the following analytical form:

$$W_f = \frac{1}{4}x_1 + \frac{109}{12}x_2 - \frac{1}{3} \rightarrow \max,$$

$$\Leftrightarrow \Omega_f^{Ox_1x_2} : \begin{cases} -3x_1 + 11x_2 \leq 32, \\ x_2 \geq 1, \\ 9x_1 + 19x_2 \leq 28, \\ 9x_1 + 23x_2 \leq 32, \\ x_1 \geq 0, x_2 \geq 0. \end{cases}$$

The graphical solution is given on Fig. 4.




Fig. 4. Projection onto Ox_1x_2

The solution of the system is the optimum vertex

$$X_{\max}^{opt} : \omega_2 \times \omega_4 \Leftrightarrow \begin{cases} x_1 = 0, \\ 9x_1 + 23x_2 = 32, \end{cases} \Leftrightarrow \begin{cases} x_1 = 0, \\ x_2 = \frac{32}{23}. \end{cases}$$

We obtain the other coordinates from the solved system (6) The problem solution is equal to:

$X_{\max}^{\text{opt}} = \left[0, \frac{32}{23}, \frac{6}{23}, \frac{3}{23}, \frac{32}{23}, 0 \right]$. The solution obtained completely agrees with the one obtained previously, with projecting onto plane Ox_3x_4 .

The biggest target function value is $W_I^{\max} = \frac{289}{23}$.

The graphic representation shows that (1,1) is the only integer point. With account taken of the previous and the current projecting, we obtain the integer problem solution $X_{\max}^Z = [1, 1, 0, 0]$. In this problem, the binary solution agrees with the integer solution $X_{\max}^{0v1} = [1, 1, 0, 0]$.

Projection onto Ox_1x_6

We project Ω_I onto Ox_1x_6 . In view of this, we take the following quadruple as basis variables: x_2, x_3, x_4, x_5 . We make calculations with the Gauss-Jordan method. (Table 3).

Table 3

	x_1	x_2	x_3	x_4	x_5	x_6	b	Σ
	2	3	4	3	1	0	7	20
	3	5	2	4	0	0	8	22
	3	3	5	4	0	0	6	21
	3	5	3	2	0	1	8	22
W_I	1	8	4	1	0	0	0	
	-4	-7	0	-5	1	0	-9	-24
	1,5	2,5	1	2	0	0	4	11
	-4,5	-9,5	0	-6	0	0	-14	-34
	-1,5	-2,5	0	-4	0	1	-4	-11
W_I	-5	-2	0	-7	0	0	-16	
	- 1/4	11/12	0	0	1	0	8/3	13/3
	0	- 2/3	1	0	0	0	- 2/3	- 1/3
	3/4	19/12	0	1	0	0	7/3	17/3
	3/2	23/6	0	0	0	1	16/3	35/3
W_I	1/4	109/12	0	0	0	0	1/3	
	- 14/23	0	0	0	1	- 11/46	32/23	71/46
	6/23	0	1	0	0	4/23	6/23	39/23
	3/23	0	0	1	0	- 19/46	3/23	39/46
	9/23	1	0	0	0	6/23	32/23	70/23
W_I	- 76/23	0	0	0	0	- 109/46	- 283/23	

From the last step of Table 3, we have the solved system

$$\begin{aligned}
 &W_I = x_1 + 8x_2 + 4x_3 + x_4 \rightarrow \max, \\
 &\Omega_I : \begin{cases} -\frac{14}{23}x_1 - \frac{11}{23}x_6 + x_5 = \frac{32}{23}, \\ \frac{6}{23}x_1 + \frac{4}{23}x_6 + x_3 = \frac{6}{23}, \\ \frac{3}{23}x_1 - \frac{19}{46}x_6 + x_4 = \frac{3}{23}, \\ \frac{9}{23}x_1 + \frac{6}{23}x_6 + x_2 = \frac{32}{23}, \\ x_1 \geq 0, x_6 \geq 0. \end{cases} \tag{7}
 \end{aligned}$$

Truncating the basis variables, we assure the transition $R^6 \Rightarrow R^2$ to two-dimensional inequalities. The

projection of six-dimensional problem onto coordinate plane Ox_1x_6 has the following form:

$$W_I = \frac{76}{23}x_1 - \frac{109}{46}x_6 + \frac{283}{23} \rightarrow \max,$$

$$\Omega_I : \begin{cases} -\frac{14}{23}x_1 - \frac{11}{46}x_6 + x_5 = \frac{32}{23}, \\ \frac{6}{23}x_1 + \frac{4}{23}x_6 + x_3 = \frac{6}{23}, \\ \frac{3}{23}x_1 - \frac{19}{46}x_6 + x_4 = \frac{3}{23}, \\ \frac{9}{23}x_1 + \frac{6}{23}x_6 + x_2 = \frac{32}{23}, \\ x_1 \geq 0, x_6 \geq 0, \end{cases} \Leftrightarrow \Omega_I^{Ox_1x_6} : \begin{cases} -28x_1 - 11x_6 \leq 64, \\ 3x_1 + 2x_6 \leq 3, \\ 6x_1 - 19x_6 \leq 6, \\ 9x_1 + 6x_6 \leq 32, \\ x_1 \geq 0, x_6 \geq 0. \end{cases}$$

The graphical solution is given on Fig. 5.




Fig. 5. Projection onto Ox_1x_6

The optimum solution is in the point of origin of coordinates

$$X_{\max}^{opt} = [0,0].$$

The other coordinates can be obtained from the solved system (7) We have:

$$X_{\max}^{opt} = \left[0, \frac{32}{23}, \frac{6}{23}, \frac{3}{23}, \frac{32}{23}, 0 \right].$$

The solution obtained is equal to the ones obtained previously, projections onto plane Ox_3x_4 and Ox_1x_2 .

The biggest value of the target function is

$$W_I^{\max} = \frac{289}{23}.$$

algorithm of such reduction proposed. This step has been realized in the environment of the Maple[®] symbolic mathematics software package. A program has been developed that provides automation of calculations using the procedure proposed. The program includes two units:

- selection of a basis variables combination and solution of the system of constraints with the Gauss-Jordan method;
- three-level optimization calculation ($x_j \geq 0, x_j \geq 0, x_j \geq 0$ and integers, $x_j = 0 \vee 1, j = 1, 2, \dots, n$) with using the standard subprogram library.

A program code fragment is given below.

The software implementation of a linear optimization problem reduction is an important component of the

```

#
eq1:=2*x[1]+3*x[2]+4*x[3]+3*x[4]+x[5]=7:
eq2:=3*x[1]+5*x[2]+2*x[3]+4*x[4]=8:
eq3:=3*x[1]+3*x[2]+5*x[3]+4*x[4]=6:
eq4:=3*x[1]+5*x[2]+3*x[3]+2*x[4]+x[6]=8:
#####

#####
W[I]=zf-max;
eq1;eq2;eq3;eq4;
x[j]>=0;

ww1:=solve({eq1,eq2,eq3,eq4},{x[3],x[4],x[5],x[6]});

om1:=-coeff(rhs(ww1[1]),x[1])*x[1]-coeff(rhs(ww1[1]),x[2])*x[2]<=
rhs(ww1[1])+(-coeff(rhs(ww1[1]),x[1])*x[1]-
coeff(rhs(ww1[1]),x[2])*x[2]):

om2:=-coeff(rhs(ww1[2]),x[1])*x[1]-coeff(rhs(ww1[2]),x[2])*x[2]<=
rhs(ww1[2])+(-coeff(rhs(ww1[2]),x[1])*x[1]-
coeff(rhs(ww1[2]),x[2])*x[2]):

om3:=-coeff(rhs(ww1[3]),x[1])*x[1]-coeff(rhs(ww1[3]),x[2])*x[2]<=
rhs(ww1[3])+(-coeff(rhs(ww1[3]),x[1])*x[1]-
coeff(rhs(ww1[3]),x[2])*x[2]):

om4:=-coeff(rhs(ww1[4]),x[1])*x[1]-coeff(rhs(ww1[4]),x[2])*x[2]<=
rhs(ww1[4])+(-coeff(rhs(ww1[4]),x[1])*x[1]-
coeff(rhs(ww1[4]),x[2])*x[2]):

zf:=subs({x[3]=rhs(ww1[1]),x[4]=rhs(ww1[2])},zf):

W[I]=zf-max;
om1;om2;om3;om4;
sort(x[1]>=0),sort(x[2]>=0);

p1:=inequal([om1,om2,om3,om4,x[1]>=0,x[2]>=0],x[1]=-1..7,x[2]=-
1..7,optionexcluded=(colour=white),
optionsfeasible=(colour=green,thickness=1)):
display(p1);

```

The results obtained within the research have allowed widening the educational content of disciplines taught within the boundaries of the educational program "Project Management". For instance, the content of discipline "Mathematical Models and Methods in Project Management" includes materials from such basic areas: linear models and linear optimization, discrete optimization, elements of the game theory. All the areas require computer modelling and IT projecting. In view of this, the proposed approach of automating optimization problems calculations is used in the educational process of training students aimed at Master's degree.

Conclusions

The proposed approach to simplification of combinatorial solution of a discrete optimization problem has significant advantages over the known methods of the optimum solution determining – the simplex method or the artificial basis method. The actually performed

decomposition of the system reduces the dimension of the equation system to be solved. Projection of the multidimensional system of the output problem onto the two-dimensional coordinate plane allows obtaining a simple system of graphical solutions for a complicated linear discrete optimization problem. From the practical point of view, the approach proposed enables reduction of compliance when calculating optimization problems of such class, and the software implementation allows including this class of problems into educational projects. The scientific result obtained makes researchers arriving at the conclusion that in the general case, it is not necessary to search for solution in all the projections. It is enough to find the solution just in one projection.

The applied significance of the approach proposed consists in using the obtained result to assure the possibility of improving complicated systems described by linear equation systems with linear constraint systems

included. Multivalued combinatorial projections cause the possibility of changing the range of problem parameters. The research has proposed projecting a multidimensional optimization process onto the two-dimensional plane.

Such method of simplification can only be applied to adapted classes of problems. The m rank of the matrix of factors of the system of constraints for a linear discrete optimization problem has to meet the condition $n-m=2$, where n – the problem dimension. It is reasonable to generalize such projecting onto three-dimensional space.

1. It has been shown that solving a linear optimization problem is possible due to simplifying through decomposition of the system by means of building projections of the multidimensional system of the output problem onto two-dimensional coordinate planes.

2. It has been confirmed on the example of solving a standard model problem that the approach proposed enables obtaining a simple system of graphical solutions of a complicated linear discrete optimization problem. The result obtained allows the researchers to arrive at the conclusion that in the general case, it is not necessary to search for the solution in all the projections. It is enough to find the solution just in one projection.

References

1. Korbut A.A., Finkelstein Y.Y. (1969). *Dyskretnoe prohammyrovanye [Discrete programming]*. Moscow: Nauka [in Russian]
2. Finkelstein Y.Y. (1976). *Priblizhennyye metody i prikladnyye zadachi diskretnogo programmirovaniya [Approximate methods and applied problems of discrete programming]*. Moscow: Nauka [in Russian]
3. Wagner H. (1973). *Osnovy issledovaniya operatsiy [Principles of Operations Research]*. Moscow: Mir
4. Burkov V. N., Gorgidze I. A., Lovetskiy S. E. (1974). *Prikladnyye zadachi teorii grafov [Applied problems of the theory of graphs]*. Tbilisi: Computation Centre of Republican Adacemy of Science of Georgian SSR
5. Sigal I. K., Ivanova A. P. (2003). *Vvedenie v prikladnoe diskretnoe programmirovanie: modeli i vychislitelnyye algoritmy [Introduction to applied discrete programming: models and calculation algorithms]*. Moscow
6. F. Nozicka, J. Guddat, H. Hollatz. (1972). *The Linear Optimization Theory*. Berlin
7. Titov S. D., Chernova L. S. (2017). *Vyshcha ta prykladna matematika [Higher and applied mathematics]*. Manual: In 2 parts, Part 1, Kharkiv, Fakt
8. Lau D. (2007). *Algebra and Discrete Mathematics 1. Basic Terms of Mathematics, Algebraic Structures 1, Linear Algebra and Analytic Geometry, Numeric Algebra*. Second corrected and supplemented edition. Berlin: Springer
9. Jean Pierre David (2017). Low latency and division free Gauss-Jordan solver in floating point arithmetic. *Journal of Parallel and Distributed Computing*, 106, 185-193.
10. Lax Peter D. (2007). *Linear algebra and application*. New York, Wiley, 2-nd ed.
11. Yeremin I. I., Astafiev N. N. (1976). *Vvedenie v teoriyu lineynogo i vyipuklogo programmirovaniya [Introduction to the theory of linear and convex programming]*. Moscow: FIZMATLIT
12. Tytov S. D., Chernova L. S. (2016). *Teoriia vyznachnykiv: Navchalno - metodychnyi posibnyk [Theory of determinants: Training and methodological manual]*. Mykolaiv: Torubara V. V.
13. Teschl Gerald, Teschl Susanne. (2008). *Mathematics for Information Scientists. Volume 1: Discrete Mathematics and Linear Algebra*. Berlin, Springer
14. Buhir M. K. (1998). *Matematyka dlia ekonomistiv. Liniina alhebra, liniini modeli [Mathematics for economists. Linear algebra, linear models]*. Kyiv
15. Buhir M. K. (2003). *Matematyka dlia ekonomistiv [Mathematics for economists]*. Alma-matir Academia
16. Titov S. D., Chernov S. K., Chernova L. S. (2018). Reduction in Discrete Optimization Problem, 2018 IEEE 13th International Scientific and Technical Conference on Computer Sciences and Information Technologies, CSIT 2018.
17. Chernov S., Titov S., Chernova L., (...), Chernova L., Kolesnikova K. Algorithm for the simplification of solution to discrete optimization problems *Eastern-European Journal of Enterprise Technologies*.
18. Kovalev M. M. (1977). *Diskretnaya optimizatsiya [Discrete optimization]*. Minsk: Byelorussian State University